



Evaluating Uncertainty to Strengthen Epidemiologic Data for Use in Human Health Risk Assessments

**Carol J. Burns, J. Michael Wright, Jennifer B. Pierson,
Thomas F. Bateson, Igor Burstyn, Daniel A. Goldstein,
James E. Klaunig, Thomas J. Luben, Gary Mihlan,
Leonard Ritter, A. Robert Schnatter, J. Morel Symons,
and Kun Don Yi**

<http://dx.doi.org/10.1289/ehp.1308062>

Received: 24 December 2013

Accepted: 29 July 2014

Advance Publication: 31 July 2014

Evaluating Uncertainty to Strengthen Epidemiologic Data for Use in Human Health Risk Assessments

Carol J. Burns,¹ J. Michael Wright,² Jennifer B. Pierson,³ Thomas F. Bateson,⁴ Igor Burstyn,⁵ Daniel A. Goldstein,⁶ James E. Klaunig,⁷ Thomas J. Luben,⁸ Gary Mihlan,⁹ Leonard Ritter,¹⁰ A. Robert Schnatter,¹¹ J. Morel Symons,¹² and Kun Don Yi¹³

¹The Dow Chemical Company, Midland, Michigan, USA; ²US EPA/NCEA, Cincinnati, Ohio, USA; ³ILSI-HESI, Washington, DC, USA; ⁴US EPA/NCEA, Washington, DC, USA; ⁵Drexel University, Department of Environmental and Occupational Health, School of Public Health, Philadelphia, Pennsylvania, USA; ⁶Monsanto, St. Louis, Missouri, USA; ⁷Indiana University, Bloomington, Indiana, USA; ⁸US EPA/NCEA, Research Triangle Park, North Carolina, USA; ⁹formerly with Bayer CropScience, Research Triangle Park, North Carolina, USA; ¹⁰University of Guelph, Guelph, Ontario, Canada; ¹¹ExxonMobil Biomedical Sciences, Inc., Annandale, New Jersey, USA; ¹²E.I. du Pont de Nemours and Company, Newark, Delaware, USA; ¹³Syngenta Crop Protection, LLC, Greensboro, North Carolina, USA

Address correspondence to Jennifer B. Pierson, 1156 15th St NW, 2nd Floor, Washington, DC 20005 USA. Telephone: 202-659-3306 ext. 158. E-mail: jpierson@hesiglobal.org

Running Title: Evaluating Uncertainty in Epidemiology

Acknowledgments: A multidisciplinary workshop steering team and break out group organizers authored this article. The workshop was co-chaired by C.J.B and J.M.W and was organized by the Health and Environmental Sciences Institute (HESI) (a part of the International Life Sciences Institute, ILSI) as an emerging issue. The contributions of the HESI Epidemiology

Subcommittee and invited participants are greatly appreciated. Dr. Gary Mihlan passed away on September 12, 2013. Gary was a valued contributor to this project and manuscript during his time at Bayer CropScience.

Participating Organizations from the HESI Epidemiology Subcommittee include: Bayer CropScience, DLW Consulting Services, LLC, The Dow Chemical Company, Drexel University, E.I. du Pont de Nemours and Company, ExxonMobil Biomedical Sciences, Inc., Harvard School of Public Health, Indiana University, Monsanto, Procter & Gamble Co., Shell Oil Company, Syngenta Crop Protection, LLC, University of Aarhus, University of Guelph, University of Leicester, US EPA/NCEA, and Wake Forest University.

The full list of workshop participants is available online:

(<http://www.hesiglobal.org/i4a/pages/index.cfm?pageID=3641>).

Declarations: This paper has been reviewed in accordance with the peer and administrative review policies of the U.S. Environmental Protection Agency (EPA) and ILSI-HESI. The views expressed in this report are those of the authors and do not necessarily reflect the opinions and/or policies of the authors' employer(s). Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the EPA.

All non-industry participants received support for travel and lodging with the exception of local participants.

The authors declared no potential conflicts of interests with respect to the authorship and/or publication of this article. The authors received no direct financial support for the research and/or authorship of this article.

This work was conducted under the auspices of the ILSI-HESI Subcommittee on Evaluating Causality in Epidemiologic Studies. HESI is a public, non-profit foundation whose mission is to

engage scientists from academia, government and industry to identify and resolve global health and environmental issues. HESI receives support primarily from its industry sponsors.

The opinions expressed herein are those of the authors and do not necessarily represent the views of HESI.

Abstract

Background: There is a recognized need to improve the application of epidemiologic data in human health risk assessment especially for understanding and characterizing risks from environmental and occupational exposures. While there is uncertainty associated with the results of most epidemiologic studies, techniques exist to characterize uncertainty that can be applied to improve weight-of-evidence evaluations and risk characterization efforts.

Methods: This report derives from a Health and Environmental Sciences Institute (HESI) workshop held in Research Triangle Park, North Carolina, to discuss the utility of using epidemiologic data in risk assessments, including the use of advanced analytical methods to address sources of uncertainty. Epidemiologists, toxicologists, and risk assessors from academia, government and industry convened to discuss uncertainty, exposure assessment, and application of analytical methods to address these challenges.

Synthesis: Several recommendations emerged to help improve the utility of epidemiologic data in risk assessment. For example, improved characterization of uncertainty is needed to allow risk assessors to quantitatively assess potential sources of bias. Data are needed to facilitate this quantitative analysis, and interdisciplinary approaches will help ensure sufficient information is collected for a thorough uncertainty evaluation. Advanced analytical methods and tools such as directed-acyclic graphs (DAGs) and Bayesian statistical techniques, can provide important insights and support interpretation of epidemiologic analysis.

Conclusions: The discussions and recommendations from this workshop demonstrate that there are practical steps that the scientific community can adopt to strengthen epidemiologic data for decision making.

Background and Introduction

Human health risk assessments have traditionally relied heavily on toxicological and other experimental data, but there is an increased recognition of the value of using epidemiologic data in risk assessment. Previous publications (Fann et al. 2011; Jones et al. 2009; Lavelle et al. 2012; Vlaanderen et al. 2008) and initiatives have discussed how to improve the application of these epidemiologic data to risk assessments. As an example, in early 2010, the U.S. EPA requested input from the Federal Insecticide, Fungicide and Rodenticide Act Scientific Advisory Panel on approaches for the “[i]ncorporation of epidemiology and human incident data into risk assessment[s].” (U.S. EPA 2009a). Epidemiologic studies play a key role in setting national ambient air quality standards (U.S. EPA 2009b) and contribute substantially to other thematic weight-of-evidence approaches toward evaluating causality based on multiple lines of evidence (Rhomborg et al. 2010; Weed 2005).

The incorporation of epidemiologic evidence into risk assessments is an important part of understanding and characterizing risks from environmental and occupational exposures. Uncertainty arises from study limitations regarding internal validity, including exposure assessment, confounding and other potential sources of bias, and external validity or generalization from study populations to the populations for which risk assessments are conducted (Guzelian et al. 2005; Hertz-Picciotto 1995; Lash et al. 2009; Levy 2008; Maldonado 2008; Persad and Cooper 2008). Further, point estimates can be inaccurate because of internal validity issues and since confidence intervals only focus on the potential for random error. These different sources of uncertainty can have an impact on various steps of the risk assessment paradigm (including hazard identification, exposure assessment and dose-response assessment)

in that hazards are not recognized, hazards are incorrectly identified, or inaccurate dose-response characterizations lead to over- or under-estimation of “safe” exposure levels.

Epidemiologic approaches and statistical techniques exist to characterize uncertainty that can be applied to weight-of-evidence evaluations and risk characterization efforts. While there is strong theoretical support for the utility of these approaches, their translation into regular epidemiologic practice is lagging. In addition, the impact of potential sources of error in epidemiologic studies is often only qualitatively discussed. For example, with respect to exposure measurement error, Jurek et al. (2006) sampled papers from three epidemiology journals over one year and found that only 61% of the articles made any mention of exposure measurement error, of which only 46% qualitatively described the possible effects. Only one of 57 sampled studies quantified the likely impact of exposure measurement error on results. This incomplete information demonstrates an opportunity among epidemiologists to characterize the magnitude and impact of various sources of uncertainty, which can help address one of the more difficult challenges in risk assessment.

This report derives from a workshop held in Research Triangle Park, North Carolina, in October 2012 to discuss the utility of using epidemiologic data in risk assessments, including the use of advanced analytical methods to address sources of uncertainty. The objective of the workshop was to develop recommendations on strengthening epidemiologic studies, so that these data can more effectively be integrated in risk assessments. The HESI workshop was focused specifically on uncertainty, exposure assessment, and application of analytical methods to address these challenges. Cross-disciplinary experts in epidemiology, toxicology, exposure assessment, and risk assessment attended the workshop. A full list of workshop participants is available online (<http://www.hesiglobal.org/i4a/pages/index.cfm?pageID=3641>). These deliberations highlighted

opportunities for epidemiologists to enhance scientific research in general, as well as to address issues related to the development and use of epidemiologic data in risk assessment.

Uncertainty

The National Research Council (NRC 2009) defined uncertainty as the “lack or incompleteness of information” critical for the risk assessment process. Uncertainty in an epidemiologic study can arise from both random and systematic error from a study, while uncertainty in a risk assessment can arise from internal and external validity concerns arising from one study or set of studies. Thus, the characterization of scientific uncertainty can provide risk assessments with a level of confidence regarding decisions that are being made and allows for evaluation of the degree that uncertainty plays in the analysis of consequences of specific policies. The NRC recommended that “risk assessments should characterize and communicate uncertainty in all key computational steps of risk assessments” while recognizing that “uncertainty analysis and characterization pose difficult technical issues, and in general related best practices have not been established.” Thus, determining the nature and magnitude of uncertainties remains one of the key challenges in risk assessment.

Because results across epidemiologic, toxicologic and clinical data may be discordant at times, there is a distinct need to understand and characterize sources of uncertainty within each of these areas to characterize potential risk and hazard for risk assessment purposes. A comprehensive analysis of uncertainty across all data sources can act as a bridge to foster the integration necessary to focus further research, improve risk assessment, and understand potential impacts on human health.

Uncertainty Issues and Recommendations

1) Improved characterization and discussion of plausible sources of uncertainty would be beneficial in all epidemiologic reports and publications

The potential for bias in epidemiologic studies is routinely acknowledged in published reports but is nearly always limited to a qualitative discussion (Jurek et al. 2006). Even quantitative discussions of selection bias, for example, are typically limited to examinations of participation rates or to potential sampling bias due to self-selection. In addition, the potential for residual confounding by measured or unmeasured factors is often acknowledged but the magnitude and direction are usually unknown or unstated. Thus, characterizing and documenting the relationships (i.e. the direction and magnitude of associations) among potential confounders, exposures and outcomes of interest is critical. Knowing the direction of a potential confounder (e.g., positive or negative confounding) could enable epidemiologic data to be used in the hazard identification stage of a risk assessment or for dose-response assessments if the magnitude of confounding is also known or the uncertainty from this source of bias could be quantified. Addressing these possible sources of bias in an epidemiologic study may allow risk assessors, to the extent possible, to quantify the consequences of any bias in a specific study or across a group of similar studies. Although not a type of bias, an additional source of uncertainty is related to generalizing study results beyond the sample population being examined in an epidemiologic study. Characterizing variability in risk among different susceptible populations will ultimately make epidemiologic study results more relevant to risk assessment efforts and risk management decision-making.

2) Conduct more validation studies and uncertainty analyses of epidemiologic study findings

The overall impact of different sources of uncertainty on epidemiologic results is infrequently considered in epidemiologic publications and data sufficient to allow the reader to undertake independent uncertainty assessments are often not presented (Jurek et al. 2007). This is essentially the lowest tier (i.e., Tier 0) of uncertainty analyses recognized by the National Research Council (2009):

- Tier 0: Default assumptions—single value of result.
- Tier 1: Qualitative but systematic identification and characterization of uncertainty.
- Tier 2: Quantitative evaluation of uncertainty making use of bounding values, interval analysis, and sensitivity analysis.
- Tier 3: Probabilistic assessment with single or multiple outcome distributions reflecting uncertainty and variability.

It is, therefore, recommended that investigators obtain additional data needed to facilitate uncertainty analysis and undertake at least a qualitative assessment of uncertainty, including all recognized sources or justifying their omission. A qualitative characterization would include identifying possible sources and beginning to assess the sources, direction and magnitude of uncertainty. The potential ramifications of each source of uncertainty should be addressed and some crude classification or categorization approaches could be developed (e.g. low, intermediate or high uncertainty) with respect to a given source. When sources of uncertainty can be identified but not fully quantified within a study or set of studies, there may be default data available that can be useful in estimating a possible range of values (Sturmer et al. 2007). Indeed, there may also be a complete lack of data that contribute to the uncertainty. However, investigators could consider the direction and magnitude of the potential uncertainty (i.e. confounding and/or bias). Such data would allow for higher tiers of uncertainty analyses.

Methodologic guidance and software for quantitative bias analysis have also become available (Lash et al. 2009) but are not yet common in risk assessment. Ideally, to facilitate the highest tier of uncertainty analysis, a quantitative assessment of individual and conjoint sources of uncertainty would be included in every epidemiologic study. The conduct of more validation studies and sensitivity analyses is also recommended to better understand methodological issues and sources of uncertainty (Chatterjee and Wacholder 2002), (e.g. Greenland 1996; Rosenbaum 2005; Schneeweiss 2006; VanderWeele and Arah 2011).

3) Improve communication about epidemiologic uncertainty

Full disclosure of uncertainty in epidemiology is encouraged as a matter of transparency. Characterization and quantification of uncertainty should increase such that the basis of decisions and assumptions are clear, either within the publication or in supplemental information. Epidemiologists and their peer scientists should encourage publications and other communications to include the necessary study-specific data on internal data relationships relevant to selection bias, information bias and confounding for quantitative bias analysis to assess uncertainty (Lash et al. 2009). Reviewers of manuscripts should also recommend qualitative, and if possible quantitative, discussions of uncertainties. The objective is that such information be more routinely collected and reported.

4) Develop a broader matrix of sources of uncertainty for the overall risk assessment process, with the goal of harmonization of uncertainty assessment across different disciplines

Risk assessments consider the totality of the evidence (i.e., epidemiology, toxicology, and other lines of scientific evidence, as well as any other knowledge in the context of risk) when

determining the weight of evidence. Other lines of evidence such as toxicology and mode of action can be used to inform the interpretation and use of epidemiologic data in risk assessment. Since uncertainties exist in all lines of scientific endeavor, each source of uncertainty across these areas should be considered in assessing uncertainty in the overall risk assessment. Previous efforts have recommended harmonization of the incorporation of uncertainty in risk assessment, primarily focusing on the use of default uncertainty values from toxicological data (Sonich-Mullin et al. 2001). Consequently, harmonization of sources of uncertainty across epidemiology and toxicology should be undertaken in a systematic manner that will make for more transparent decision-making.

Exposure Assessment

Exposure science aims to quantify the intensity, frequency, duration, and timing of human contact with chemical, physical, or biological agents occurring in the environment, and may be used to further inform evaluation of causality in the environmental source-to-health outcome continuum (Barr 2006). Within exposure science, exposure assessment specifically deals with several distinct aspects that underlie the risk assessment process, including the exposure source(s), the environmental pathway(s), environmental concentrations, human exposures, and dose.

Data are rarely available on biologically relevant dose metrics (e.g., absorbed dose, effective dose, etc.) in the organ or tissue of interest in epidemiologic-studies; thus, dose is often estimated indirectly using exposure metrics. These surrogate estimates of exposure are subject to measurement error since they may rely on imperfectly measured concentrations in the individual, or models of transport and fate in the environment or workplace. Additionally, measurement

error may result from estimates of the distribution of human uptake over time (e.g., use of physiologically based pharmacokinetic model), or collection of activity use pattern data.

Similar to measurement error and resulting misclassification of health outcome data, exposure misclassification is important to characterize in epidemiologic studies because it can distort exposure-response relationships and lead to biased or imprecise results. Exposure measurement error can be differential or non-differential with respect to variation in disease status. Exposure measurement error can lead to exposure misclassification when exposure surrogates for individual participants are classified into categories for analysis.

Differential misclassification can arise in categorical exposure metrics even when there is non-differential error (i.e., independent of disease status) in an exposure variable that is measured on a continuous scale (Flegal et al. 1991). In order for epidemiologic studies to be evaluated and utilized appropriately in risk assessment, it is important that exposure measurement error is characterized and evaluated thoroughly with consideration of the magnitude and direction of any potential exposure misclassification bias (for example, see Bergen et al. 2013). This information is useful for risk assessors when they evaluate the potential for bias and confidence placed on study results.

Exposure Assessment Issues and Recommendations

1) An interdisciplinary perspective is needed during the study design phase to ensure that biologically relevant quantitative exposure-response information is collected that will be useful for risk assessment purposes

During the study-design phase, an interdisciplinary team including experts, for example, in epidemiology, exposure assessment, industrial hygiene, and analytical chemistry, should be

assembled to develop robust exposure assessment approaches. This might include consideration of targeted data collection strategies, such as collection of exposure or surrogate data based on the appropriate biological matrix, sample number, and the critical exposure window(s). Other constraints that can be addressed include sources of exposure variability, availability of resources, participant burden and ethical considerations (with Institutional Review Board review, as appropriate). This interdisciplinary approach will allow for collection of biologically relevant exposure data to increase the potential for quantification of exposure-response relationships that will be useful for risk assessment and risk management purposes.

2) Develop exposure assessment approaches that are transparent and well-characterized

It is recommended that study authors should discuss the nature (i.e., type, direction and magnitude) and likelihood of any expected exposure measurement error and misclassification bias. An evaluation of measurement error and any resulting impact on effect estimates would provide risk assessors with information to weight studies by the quality of the exposure assessment, the methods used to adjust for exposure measurement error and the likelihood that exposure measurement error contributes to uncertainty in effect estimates in epidemiologic study results. Characterization of exposure data quality may include steps to make exposure data publicly available so that risk assessors can perform secondary data analyses, including sensitivity and uncertainty analyses.

3) Quantify exposure measurement error and examine and correct for its impact on effect estimates

Ignoring uncertainty in exposure estimation can produce bias when such estimates are used to examine associations with adverse health effects (Carroll et al. 1995). Although epidemiologic

publications infrequently present detailed information on the potential impact of measurement error (Jurek et al. 2006; Spiegelman 2010), epidemiologic study results would be enhanced by detailing exposure assessment assumptions and characterizing measurement error to allow risk assessors to gauge the potential impact of this error. This should include characterization of different sources and types of measurement error. The sources and types can be based on various assumptions used in exposure modeling efforts including unaccounted inter- and intra-individual variability in exposure patterns (Kromhout et al. 1993; Symanski et al. 2007), or from variability based on limited monitoring data. Once these different types and sources of measurement error are identified, bias analyses should be included to examine uncertainty due to the use of different exposure metrics in relation to what is known about the critical exposure period or evaluation of specific parameter estimates (e.g., half-life considerations of biological measures) or other modeling assumptions, such as the validity of the underlying input data (e.g., chemical monitoring data) and modeling data (e.g., fate and transport models) used to estimate exposure concentrations. Statistical techniques, both non-Bayesian and Bayesian, are available to allow for the correction of biased effect estimates due to exposure measurement error. Examples of non-Bayesian methods for accounting and adjusting for exposure measurement error include conditional likelihood methods (Guolo and Brazzale 2008; Lash and Fink 2003; Lash et al. 2009; Maldonado 2008; Stram et al. 2003) such as regression calibration (Spiegelman 2010) and conditional scores procedures (McShane et al. 2001), while Bayesian methods exist that can be used for both binary and continuous exposures (Espino-Hernandez et al. 2011; Liu et al. 2009; Prescott and Garthwaite 2005; Rice 2003).

4) Develop improved methods for assessing exposures to multiple environmental chemicals and multiple routes of exposure

Traditionally, risk assessments have focused primarily on single chemicals. However, this does not reflect human exposure conditions. There is a recognized need to focus on multi-media sources of exposure to individual chemicals, as well as complex mixtures. This is an area of research where observational studies, such as epidemiology, are an improvement over experimental studies because they can more readily address multiple exposures simultaneously.

It is important that epidemiologists continue to develop and evaluate methods for assessing exposure to complex mixtures in order to better characterize exposure assessment and to allow for the evaluation of effect measure modification and confounding. This would establish a robust scientific database necessary to conduct cumulative risk assessments. The understanding of the relationships among complex exposures will require modeling of monitoring data and other exposure determinants and development of techniques for assessing exposures to mixtures that result in unbiased or minimally biased effect estimates (Carlin et al. 2013). Additionally, approaches such as multivariate source receptor modeling represent promising avenues for assessing exposure to complex mixtures (Hopke 2010), although further work is needed to account for key sources of uncertainty in such models. The development of efficient, easily-measured, cost-effective exposure surrogates for key mixtures of concern will be important, and will include its own challenges for identifying and quantifying exposure measurement error. For example, it will be important to understand how the type and structure of measurement error may differ across different individual mixture components or for a surrogate representing exposure to the whole mixture. Techniques are needed to characterize and adjust for exposure measurement error of chemical mixtures.

Analytic Tools in Epidemiology

As epidemiologic data are key inputs for risk assessments, it is important to apply methodologies that better characterize validity and precision of study results. The methods considered can be broadly classified as (1) frequentist methods to address study biases systematically and quantitatively, (2) Bayesian statistical techniques, which utilize prior knowledge addressing causal hypotheses and estimation problems under evaluation, and (3) computational methods (e.g., cross-validation, re-sampling techniques, and boosting and model ensemble techniques), which provide valid statistical inferences without requiring strong *a priori* modeling assumptions. Each of these broad approaches addresses validity and characterizes the uncertainty of results from a single study and extends to improved characterization of epidemiologic results in weight of evidence assessments.

The analytic methods group discussion included four specific areas that facilitate causal interpretation in epidemiology: (1) the use of directed acyclic graphs [DAGs - diagrams consisting of variables connected by arrows or lines to depict often complex relationships (Joffe et al. 2012)], (2) summarizing epidemiologic results using Bayesian posterior distributions, (3) strategies for quantitatively evaluating measurement error, and (4) formally assessing causality as it relates to policy decisions. Each of these areas led to a set of recommendations. These areas served as the basis for further discussion of related topics such as primary versus secondary analyses, journal requirements, epidemiology curricula, and data sharing practices.

Analytic Methods: Issues and Recommendations

1) The application of directed acyclic graphs (DAGs) should be encouraged more broadly

Joffe and colleagues (2012) describe how DAGs make explicit the assumed or estimated relationships among unobserved and measured variables, indicating the causal direction of the potential relationships. As described, DAGs are considered to be an appropriate method to illustrate causal hypotheses and to specify the structure of associations between variables of interest. They also provide a useful way to represent assumptions, especially conditional independence assumptions, necessary for statistical analyses and causal inference. Lastly, DAGs are helpful for determining which factors may be confounders or effect modifiers of an association between exposure and outcome (VanderWeele and Robins 2007). DAGs provide transparent representations of a hypothesis as well as justification for specific analytic strategies to be applied during the investigation such as identification of causal intermediates. DAGs can also clarify methodologic challenges, such as illustrating selection bias (Flanders and Klein 2007; Hernán et al. 2004). It is recommended that journal editors request that DAGs be included in supplementary material (Westreich and Greenland 2013).

2) Incorporate prior knowledge through Bayesian methods

Bayesian statistical analysis differs from frequentist methods in that Bayesian analyses use information that exists before study data are collected and analyzed (i.e., “prior” distributions) to update what can be learned about a specific problem after conducting a study by expressing new state of knowledge as “posterior” distributions. Results from the literature or other data sources are used to specify the *a priori* distribution for any parameters, such as the size and direction of exposure-outcome associations and the extent of measurement error. Subsequently, the study

results generated by the analysis can assess the conditional probability distribution of parameters of interest (the posterior distribution) by reconciling the data observed, the analytic model fitted to the study data, and the prior information incorporated into the analytic model (Bolstad 2007).

Bayesian techniques can also allow for simultaneous correction for sources of bias such as measurement error and confounding (de Vocht et al. 2009; Steenland and Greenland 2004) that are typically treated in isolation in current practice of epidemiology (Gustafson and McCandless 2010). Although these techniques are not routinely employed, specification of prior model probabilities by investigators is inherent in grant proposals, the introduction section of a study publication, and the subjective interpretation of results (Goodman 2001). Thus, it could be argued that current practice involves presenting Bayesian considerations of a research article, while the reported results often rely on frequentist analysis and qualitative interpretations (Pearce and Corbin 2013).

3) Measurement error should not be ignored in any analysis of epidemiologic results and should be assessed using quantitative methods

Measurement error is an almost universal limitation of epidemiologic studies and their analyses. The current practice of acknowledging it diffusely with a brief discussion that frequently invokes its theoretical impact, e.g., that it is most likely to be non-differential and results in potential for bias towards a null result, will not improve epidemiologic input into risk assessments. Strategies for correcting for the bias due to measurement error quantitatively are described in textbooks and can be readily implemented for many study designs. These include regression calibration, simulation-extrapolation, Bayesian approaches (Carroll et al. 1995; Gustafson 2004), and computational statistical approaches (e.g., multiple imputations, data augmentation, and expectation-maximization algorithms). Attention should be given to correcting for measurement

error available in commonly used epidemiologic software platforms such as *rcal* in STATA (<http://www.stata.com/merror/rcal.pdf>) (Hardin 2003) or *PROC CALIS* in SAS. Peer reviewers and journal editors should expect formal quantitative assessments of measurement error and related biases, as well as correction for bias that it creates, rather than relying on qualitative discussions. In addition, adequate funding should be designated for exposure validation studies and granting agencies should allow for such validations as essential criteria for funding epidemiologic research (Heid et al. 2004).

4) Distinguish associations from causes

Formal causality assessments are important and impact policy decisions. The synthesis of epidemiologic studies can be the primary basis for regulation and policy actions. Without state-of-the-art analytic techniques being used more routinely in epidemiologic studies and other lines of evidence, the benefits and costs of recommended interventions or action could be misestimated; and apparent cost-effective interventions may be ineffective. In particular, it is unwarranted to assume that a specific statistical association represents a causal effect, such that changing the predictor variable would cause a corresponding change in the outcome variable (Freedman 2004). Indeed, the distinction between structural and reduced-form equations in econometrics, and phenomena such as Simpson's Paradox, demonstrate that (reduced-form) regression coefficients need not even have the same sign as corresponding causal coefficients showing how a change in an explanatory variable would change the dependent variable (Pearl 2009).

Although epidemiologists are aware of basic threats to inferential validity from observational studies, there is little agreement, even amongst workshop participants, on whether

epidemiologists should consider policy implications of declaring an association to be causal. One view expressed was that epidemiologists should primarily conduct research that supports or refutes qualitative statements about causation, as in showing that an exposure “causes” a specific disease. This viewpoint emphasizes epidemiology’s role in hazard identification, i.e., an early stage of risk assessment for which putative threats to health are identified as causal. Another viewpoint was that epidemiologic results could be used to conceptualize causation in the context of population health, as in showing that some modifiable exposure is capable of causing important changes in health of population overall. This would more closely align epidemiology with the risk characterization phase of risk assessment where costs and benefits of risk management interventions are weighted and risks are appraised quantitatively (Phillips 2001). Alternative outcomes analysis is an example of a technique that can provide important insights in distinguishing association from causation, and could be more routinely applied in assessing causal inference and attributable risk estimation (Jager et al. 1990; Meijster et al. 2011a; Meijster et al. 2011b; Thomsen et al. 2006). Alternative outcomes analysis allows for the conceptualization of causation in terms of causes of meaningful versus ignorable consequences assuming these can be readily differentiated into one of these two options. Regardless of how epidemiologic data align with the risk assessment paradigm, epidemiologic practice should adopt methods that apply state-of-the-art techniques to address uncertainty and other study limitations and to help contextualize epidemiologic study results in terms of causality and public health intervention.

Conclusions and Future Directions

Epidemiologic data are critical for risk assessment efforts, but are rarely conducted with quantification of uncertainty, which may limit their use in risk assessments. The HESI

Epidemiology Subcommittee workshop focused on strengthening the utility and application of epidemiology studies by recommending improvements in analytic methods, exposure assessment approaches and other techniques to quantify and account for specific sources of uncertainty.

Several recommendations resulted from this effort. Specific statistical approaches and analytical techniques, such as increased use of quantitative bias-analysis, DAGs and Bayesian analyses, are available for improving the inferences drawn from epidemiologic results and currently used infrequently. Additionally, new methods may be needed for assessing exposure and characterizing uncertainty related to chemical mixtures. Other deliberations in the workshop highlighted the complete reporting of all data elements and analytical tables to permit others to conduct uncertainty analyses (either in the journal supplemental appendices or through the investigators' institution). Specifically, increased transparency of results would improve weight of evidence evaluations and collaboration amongst researchers in other disciplines would improve study designs and analytical approaches, particularly for exposure assessment.

Although there are multiple strategies for quantifying and reducing measurement error, there are barriers for routinely applying these techniques. A key disincentive is that substantial time and effort can be required to conduct validation or reliability studies, which can put a strain on research budgets. There may also be a perception that analyses of exposure measurement error tends to decrease the estimated precision of reported results, thereby increasing the probability of a “false-negative” result (Blair et al. 2009). It has been suggested that exposure measurement error and resulting misclassification is more likely to be non-differential by disease status in epidemiology studies and will most frequently result in false negatives through attenuation of effect estimates (Blair et al. 2007). This assumption is made despite evidence from statistical literature that the impact of exposure measurement error can be profound and complex and that it

is difficult to anticipate its impact on effect estimates in an individual study (Gustafson 2004). Since many manuscripts are routinely accepted without analyses quantifying uncertainty, validated exposure assessment or use of advanced analytic methods, there is little incentive to adopt the recommendations made. Funding organizations, peer reviewers, and journal editors should be catalysts for change in this effort.

The discussions and recommendations from this workshop demonstrate that there are practical steps that the scientific community can adopt to strengthen epidemiology data for decision making. Use of available methods to quantify and adjust for uncertainty will help reduce the potential impact of different sources of error and bias and help achieve better decisions for risk assessment, policy and ultimately, public health.

References

- Barr D. 2006. Human exposure science: A field of growing importance. *J Expo Sci Environ Epidemiol* 16:473; doi: 10.1038/sj.jes.7500536.
- Bergen S, Sheppard L, Sampson PD, Kim SY, Richards M, Vedal S, et al. 2013. A national prediction model for PM2.5 component exposures and measurement error-corrected health effect inference. *Environ Health Perspect* 121:1017-1025; doi: 10.1289/ehp.1206010.
- Blair A, Saracci R, Vineis P, Cocco P, Forastiere F, Grandjean P, et al. 2009. Epidemiology, public health, and the rhetoric of false positives. *Environ Health Perspect* 117:1809-1813; doi: 10.1289/ehp.0901194.
- Blair A, Stewart P, Lubin JH, Forastiere F. 2007. Methodological issues regarding confounding and exposure misclassification in epidemiological studies of occupational exposures. *Am J Ind Med* 50:199-207; doi: 10.1002/ajim.20281.
- Bolstad WM. 2007. Introduction to bayesian statistics, 2nd Edition: John Wiley & Sons, Inc.
- Carlin DJ, Rider CV, Woychik R, Birnbaum LS. 2013. Unraveling the health effects of environmental mixtures: An NIEHS priority. *Environ Health Perspect* 121:A6-A8; doi: 10.1289/ehp.1206182.
- Carroll RJ, Ruppert D, Stefanski LA. 1995. Measurement error in nonlinear models. London: Chapman & Hall.
- Chatterjee N, Wacholder S. 2002. Validation studies: Bias, efficiency, and exposure assessment. *Epidemiology* 13:503-506.
- de Vocht F, Kromhout H, Ferro G, Boffetta P, Burstyn I. 2009. Bayesian modelling of lung cancer risk and bitumen fume exposure adjusted for unmeasured confounding by smoking. *Occup Environ Med* 66:502-508; doi: 10.1136/oem.2008.042606.
- Espino-Hernandez G, Gustafson P, Burstyn I. 2011. Bayesian adjustment for measurement error in continuous exposures in an individually matched case-control study. *BMC Med Res Methodol* 11:67; doi: 10.1186/1471-2288-11-67.
- Fann N, Bell ML, Walker K, Hubbell B. 2011. Improving the linkages between air pollution epidemiology and quantitative risk assessment. *Environ Health Perspect* 119:1671-1675; doi: 10.1289/ehp.1103780.
- Flanders WD, Klein M. 2007. Properties of 2 counterfactual effect definitions of a point exposure. *Epidemiology* 18:453-460; doi: 10.1097/01.ede.0000261472.07150.4f.

- Flegal KM, Keyl PM, Nieto FJ. 1991. Differential misclassification arising from nondifferential errors in exposure measurement. *Am J Epidemiol* 134:1233-1244.
- Freedman DA. 2004. Graphical models for causation, and the identification problem. *Eval Rev* 28:267-293; doi: 10.1177/0193841X04266432.
- Goodman SN. 2001. Of P-values and Bayes: A modest proposal. *Epidemiology* 12:295-297; doi: 10.1097/00001648-200105000-00006.
- Greenland S. 1996. Basic methods for sensitivity analysis of biases. *Int J Epidemiol* 25:1107-1116.
- Guolo A, Brazzale AR. 2008. A simulation-based comparison of techniques to correct for measurement error in matched case-control studies. *Stat Med* 27:3755-3775; doi: 10.1002/sim.3282.
- Gustafson P. 2004. Measurement error and misclassification in statistics and epidemiology: Impacts and Bayesian Adjustments. Chapman & Hall/CRC Press.
- Gustafson P, McCandless LC. 2010. Probabilistic approaches to better quantifying the results of epidemiologic studies. *Int J Environ Res Public Health* 7:1520-1539; doi: 10.3390/ijerph7041520.
- Guzelian PS, Victoroff MS, Halmes NC, James RC, Guzelian CP. 2005. Evidence-based toxicology: A comprehensive framework for causation. *Hum Exp Toxicol* 24:161-201; doi: 10.1191/0960327105ht517oa.
- Hardin JW, Carroll, R.J., Schmiediche, H. 2003. The regression calibration method for fitting generalized linear models with additive measurement error. *Stata J*:1-11.
- Heid IM, Kuchenhoff H, Miles J, Kreienbrock L, Wichmann HE. 2004. Two dimensions of measurement error: Classical and Berkson error in residential radon exposure assessment. *J Expo Anal Environ Epidemiol* 14:365-377; doi: 10.1038/sj.jea.7500332.
- Hernán MA, Hernandez-Diaz S, Robins JM. 2004. A structural approach to selection bias. *Epidemiology* 15:615-625; doi: 10.1097/01.ede.0000135174.63482.43.
- Hertz-Picciotto I. 1995. Epidemiology and quantitative risk assessment: A bridge from science to policy. *Am J Public Health* 85:484-491; doi: 10.2105/AJPH.85.4.484.
- Hopke PK. 2010. The application of receptor modeling to air quality data. *Pollution Atmosphérique* 91-109.

- Jager JC, Postma MJ, Boom FM, Reinking DP, Borleffs JCC, Heisterkamp SH, et al. 1990. Epidemiological models and socioeconomic information: Methodological aspects of AIDS/HIV scenario analysis. In: Economic aspects of AIDS and HIV infection, (Schwefel D, Leidl R, Rovira J, Drummond M, Eds):Springer Berlin Heidelberg, 262-281.
- Joffe M, Gambhir M, Chadeau-Hyam M, Vineis P. 2012. Causal diagrams in systems epidemiology. *Emerg Themes Epidemiol* 9:1; doi: 10.1186/1742-7622-9-1.
- Jones DR, Peters JL, Rushton L, Sutton AJ, Abrams KR. 2009. Interspecies extrapolation in environmental exposure standard setting: A bayesian synthesis approach. *Regul Toxicol Pharmacol* 53:217-225; doi: 10.1016/j.yrtph.2009.01.011.
- Jurek AM, Maldonado G, Greenland S, Church TR. 2006. Exposure-measurement error is frequently ignored when interpreting epidemiologic study results. *Eur J Epidemiol* 21:871-876; doi: 10.1007/s10654-006-9083-0.
- Jurek AM, Maldonado G, Greenland S, Church TR. 2007. Uncertainty analysis: An example of its application to estimating a survey proportion. *J Epidemiol Community Health* 61:650-654; doi: 10.1136/jech.2006.053660.
- Kromhout H, Symanski E, Rappaport SM. 1993. A comprehensive evaluation of within- and between-worker components of occupational exposure to chemical agents. *Ann Occup Hyg* 37:253-270; doi: 10.1093/annhyg/37.3.253.
- Lash TL, Fink AK. 2003. Semi-automated sensitivity analysis to assess systematic errors in observational data. *Epidemiology* 14:451-458; doi: 10.1097/01.EDE.0000071419.41011.cf.
- Lash TL, Fox MP, Fink AK. 2009. Applying quantitative bias analysis to epidemiologic data. Vol. XII. New York:Springer.
- Lavelle KS, Robert Schnatter A, Travis KZ, Swaen GM, Pallapies D, Money C, et al. 2012. Framework for integrating human and animal data in chemical risk assessment. *Regul Toxicol Pharmacol* 62:302-312; doi: 10.1016/j.yrtph.2011.10.009.
- Levy JI. 2008. Is epidemiology the key to cumulative risk assessment? *Risk Anal* 28:1507-1513; doi: 10.1111/j.1539-6924.2008.01121.x.
- Liu J, Gustafson P, Cherry N, Burstyn I. 2009. Bayesian analysis of a matched case-control study with expert prior information on both the misclassification of exposure and the exposure-disease association. *Stat Med* 28:3411-3423; doi: 10.1002/sim.3694.

- Maldonado G. 2008. Adjusting a relative-risk estimate for study imperfections. *J Epidemiol Community Health* 62:655-663; doi: 10.1136/jech.2007.063909.
- McShane LM, Midthune DN, Dorgan JF, Freedman LS, Carroll RJ. 2001. Covariate measurement error adjustment for matched case-control studies. *Biometrics* 57:62-73.
- Meijster T, van Duuren-Stuurman B, Heederik D, Houba R, Koningsveld E, Warren N, et al. 2011a. Cost-benefit analysis in occupational health: A comparison of intervention scenarios for occupational asthma and rhinitis among bakery workers. *Occup Environ Med* 68:739-745; doi: 10.1136/oem.2011.064709.
- Meijster T, Warren N, Heederik D, Tielemans E. 2011b. What is the best strategy to reduce the burden of occupational asthma and allergy in bakers? *Occup Environ Med* 68:176-182; doi: 10.1136/oem.2009.053611.
- NRC. 2009. Science and decisions: Advancing risk assessment: The National Academies Press.
- Pearce N, Corbin M. 2013. Why should we be bayesians (and often already are without realizing it). In: *Current topics in occupational epidemiology* (Venableskm, Ed) Oxford, England: Oxford University Press, 218-233.
- Pearl J. 2009. Chapter 7 In: "Causality: Models, reasoning and inference. In: *The logic of structure-based counterfactuals*, 2nd Edition: Cambridge University Press.
- Persad AS, Cooper GS. 2008. Use of epidemiologic data in integrated risk information system (iris) assessments. *Toxicol Appl Pharmacol* 233:137-145; doi: 10.1016/j.taap.2008.01.013.
- Phillips CV. 2001. The economics of 'more research is needed'. *Int J Epidemiol* 30:771-776; doi: 10.1093/ije/30.4.771.
- Prescott GJ, Garthwaite PH. 2005. Bayesian analysis of misclassified binary data from a matched case-control study with a validation sub-study. *Stat Med* 24:379-401; doi: 10.1002/sim.2000.
- Rhomberg LR, Bailey LA, Goodman JE. 2010. Hypothesis-based weight of evidence: A tool for evaluating and communicating uncertainties and inconsistencies in the large body of evidence in proposing a carcinogenic mode of action - naphthalene as an example. *Crit Rev Toxicol* 40:671-696; doi: 10.3109/10408444.2010.499504.
- Rice K. 2003. Full-likelihood approaches to misclassification of a binary exposure in matched case-control studies. *Stat Med* 22:3177-3194; doi: 10.1002/sim.1546.

- Rosenbaum PR. 2005. Sensitivity Analysis in Observational Studies, Volume 4, In Encyclopedia of Statistics in Behavioral Science (Everitt BS, Howell DC Eds). John Wiley & Sons, Ltd, Chichester, 1809-1814.
- Schneeweiss S. 2006. Sensitivity analysis and external adjustment for unmeasured confounders in epidemiologic database studies of therapeutics. *Pharmacoepidemiol Drug Saf* 15:291-303; doi: 10.1002/pds.1200.
- Sonich-Mullin C, Fielder R, Wiltse J, Baetcke K, Dempsey J, Fenner-Crisp P, et al. 2001. IPCS conceptual framework for evaluating a mode of action for chemical carcinogenesis. *Regul Toxicol Pharmacol* 34:146-152; doi: 10.1006/rtph.2001.1493.
- Spiegelman D. 2010. Approaches to uncertainty in exposure assessment in environmental epidemiology. *Annu Rev Public Health* 31:149-163; doi: 10.1146/annurev.publhealth.012809.103720.
- Steenland K, Greenland S. 2004. Monte Carlo sensitivity analysis and bayesian analysis of smoking as an unmeasured confounder in a study of silica and lung cancer. *Am J Epidemiol* 160:384-392; doi: 10.1093/aje/kwh211.
- Stram DO, Leigh Pearce C, Bretsky P, Freedman M, Hirschhorn JN, Altshuler D, et al. 2003. Modeling and E-M estimation of haplotype-specific relative risks from genotype data for a case-control study of unrelated individuals. *Hum Hered* 55:179-190; doi: 10.1159/000073202.
- Sturmer T, Glynn RJ, Rothman KJ, Avorn J, Schneeweiss S. 2007. Adjustments for unmeasured confounders in pharmacoepidemiologic database studies using external information. *Med Care* 45:S158-165; doi: 10.1097/MLR.0b013e318070c045.
- Symanski E, Greeson NMH, Chan W. 2007. Evaluating measurement error in estimates of worker exposure assessed in parallel by personal and biological monitoring. *Am J Ind Med* 50:112-121; doi: 10.1002/ajim.20422.
- Thomsen M, Sorensen P, Fauser P, Porragas G. Risk scenario analysis. ISEE/ISEA 2006 conference abstracts supplement. In: Proceedings of the ISEE/ISEA 2006 Conference, 2006, Vol. 17, S486-S487.
- U.S. EPA. 2009a. FIFRA scientific advisory panel: Notice of public meeting (epa-hq-opp-2009-0851).U.S. EPA.

- U.S. EPA. 2009b. Integrated science assessment for particulate matter (final report). EPA/600/R-08/139F. Washington, D.C.
- VanderWeele TJ, Arah OA. 2011. Bias formulas for sensitivity analysis of unmeasured confounding for general outcomes, treatments, and confounders. *Epidemiology* 22:42-52; doi: 10.1097/EDE.0b013e3181f74493.
- VanderWeele TJ, Robins JM. 2007. Four types of effect modification: A classification based on directed acyclic graphs. *Epidemiology* 18:561-568.; doi: 10.1097/EDE.0b013e318127181b.
- Vlaanderen J, Vermeulen R, Heederik D, Kromhout H. 2008. Guidelines to evaluate human observational studies for quantitative risk assessment. *Environ Health Perspect* 116:1700-1705; doi: 10.1289/ehp.11530.
- Weed DL. 2005. Weight of evidence: A review of concept and methods. *Risk Anal* 25:1545-1557; doi: 10.1111/j.1539-6924.2005.00699.x.
- Westreich D, Greenland S. 2013. The table 2 fallacy: Presenting and interpreting confounder and modifier coefficients. *Am J Epidemiol* 177:292-298; doi: 10.1093/aje/kws412.